Table 7.3
Reported Deaths and Annual Death Rates Per 1,000 Patient-Years at Risk, 1998 to 2007
Pancreas After Kidney (PAK) Waiting List

Age This Year			Year										
		1998	1999	2000	2001	2002	2003	2004	2005	2006	2007		
Total													
	Patients	315	469	770	1,026	1,218	1,324	1,490	1,432	1,400	1,345		
	Deaths	5	5	10	16	18	16	26	24	30	20		
	Rate	31.3	25.0	26.4	28.1	24.9	19.1	27.7	24.5	31.0	21.6		
6-11 Years													
	Patients	-	-	-	-	-	1	-	-	_	-		
	Deaths	-	-	-	-	-	-	-	-	-	-		
	Rate	-	-	-	-	-	*	-	-	-	-		
12-17 Years													
	Patients	-	-	-	-	-	-	1	1	-	-		
	Deaths	-	-	-	-	-	-	-	-	-	-		
	Rate	-	-	-	-	-	-	*	*	-	-		
18-34 Years													
	Patients	48	96	141	183	204	198	233	200	193	153		
	Deaths	-	1	-	3	2	2	2	3	4	2		
	Rate	-	26.6	-	31.2	17.3	15.7	15.1	22.1	30.9	18.5		
35-49 Years													
	Patients	230	313	507	669	780	870	930	866	828	792		
	Deaths	5	4	9	11	9	11	17	11	21	11		
	Rate	41.9	29.5	35.7	29.6	19.2	20.1	28.5	18.5	36.6	19.9		
50-64 Years													
	Patients	37	60	122	171	230	251	321	356	372	392		
	Deaths	-	-	1	2	7	3	7	10	5	6		
	Rate	-	-	16.7	20.1	50.5	18.6	34.2	41.1	19.3	23.0		
65+ Years													
	Patients	-	-	-	3	4	4	5	9	7	8		
	Deaths	-	-	-	-	-	-	-	-	-	1		
	Rate	-	-	-	*	*	*	*	*	*	*		

Patients alive on the waiting list at any time during the year are counted. Period at risk begins the later of January 1 or waiting list registration and ends on the earlier of December 31, date of death, or date of removal for other reasons. Please see Technical Notes for further details about death rate computation.

^{(-) =} None in category.

^{(*) =} For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Table 7.3 (Continued) Reported Deaths and Annual Death Rates Per 1,000 Patient-Years at Risk, 1998 to 2007 Pancreas After Kidney (PAK) Waiting List

Ethnicity/Race			Year										
		1998	1999	2000	2001	2002	2003	2004	2005	2006	2007		
Total													
	Patients	315	469	770	1,026	1,218	1,324	1,490	1,432	1,400	1,345		
	Deaths	5	5	10	16	18	16	26	24	30	20		
	Rate	31.3	25.0	26.4	28.1	24.9	19.1	27.7	24.5	31.0	21.6		
White													
	Patients	289	415	664	872	1,040	1,118	1,249	1,171	1,150	1,088		
	Deaths	4	5	8	13	15	12	21	19	19	19		
	Rate	27.3	28.2	25.0	27.1	24.5	16.9	26.6	23.9	23.9	25.2		
African Am	nerican												
	Patients	20	33	64	82	96	107	126	151	146	145		
	Deaths	-	-	1	2	3	2	3	4	7	1		
	Rate	-	-	28.2	38.5	50.5	30.1	37.8	38.2	70.9	10.1		
Other/Mult	:i-race												
	Patients	1	4	3	7	7	8	6	9	8	6		
	Deaths	-	-	-	-	-	-	-	1	1			
	Rate	*	*	*	*	*	*	*	*	*	,		
Asian													
	Patients	2	3	5	11	9	10	10	12	11	15		
	Deaths	1	-	-	-	-	-	-	-	-	-		
	Rate	*	*	*	-	*	-	-	-	-	-		
Hispanic/L	atino.												
•	Patients	3	14	34	54	66	81	99	89	85	91		
	Deaths	-	-	1	1	-	2	2	-	3	-		
	Rate	*	-	54.5	36.0	-	41.7	34.6	-	48.8			

Patients alive on the waiting list at any time during the year are counted. Period at risk begins the later of January 1 or waiting list registration and ends on the earlier of December 31, date of death, or date of removal for other reasons. Please see Technical Notes for further details about death rate computation.

^{(-) =} None in category

^{(*) =} For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Table 7.3 (Continued)
Reported Deaths and Annual Death Rates Per 1,000 Patient-Years at Risk, 1998 to 2007
Pancreas After Kidney (PAK) Waiting List

Gender		Year									
		1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
Total											
	Patients	315	469	770	1,026	1,218	1,324	1,490	1,432	1,400	1,345
	Deaths	5	5	10	16	18	16	26	24	30	20
	Rate	31.3	25.0	26.4	28.1	24.9	19.1	27.7	24.5	31.0	21.6
Female											
	Patients	137	203	340	438	514	569	656	653	627	588
	Deaths	2	1	3	7	7	9	11	11	14	7
	Rate	29.8	11.7	18.1	29.0	22.6	24.9	26.7	25.0	32.4	17.3
Male											
	Patients	178	266	430	588	704	755	834	779	773	757
	Deaths	3	4	7	9	11	7	15	13	16	13
	Rate	32.4	35.0	32.8	27.4	26.5	14.6	28.6	24.1	29.8	24.9

Patients alive on the waiting list at any time during the year are counted. Period at risk begins the later of January 1 or waiting list registration and ends on the earlier of December 31, date of death, or date of removal for other reasons. Please see Technical Notes for further details about death rate computation.

^{(-) =} None in category.

^{(*) =} For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Table 7.3 (Continued) Reported Deaths and Annual Death Rates Per 1,000 Patient-Years at Risk, 1998 to 2007 Pancreas After Kidney (PAK) Waiting List

Blood Type						Yea	ar				
		1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
Total											
	Patients	315	469	770	1,026	1,218	1,324	1,490	1,432	1,400	1,345
	Deaths	5	5	10	16	18	16	26	24	30	20
	Rate	31.3	25.0	26.4	28.1	24.9	19.1	27.7	24.5	31.0	21.6
A											
	Patients	135	188	318	412	464	491	562	536	529	496
	Deaths	2	1	2	8	7	4	11	5	10	9
	Rate	33.6	12.8	12.7	36.4	26.0	13.2	32.6	13.9	27.7	27.4
AB											
	Patients	13	17	30	38	44	45	51	47	49	48
	Deaths	-	-	-	1	1	-	-	1	-	2
	Rate	-	-	-	45.5	43.7	-	-	34.1	-	63.6
В											
	Patients	33	56	84	112	147	161	171	144	139	137
	Deaths	-	-	1	-	1	3	3	4	5	1
	Rate	-	-	26.6	-	11.6	29.0	27.4	39.4	54.0	10.4
0											
	Patients	134	208	338	464	563	627	706	705	683	664
	Deaths	3	4	7	7	9	9	12	14	15	8
	Rate	39.2	42.0	40.8	26.3	26.0	22.2	25.9	28.6	30.9	17.0

Patients alive on the waiting list at any time during the year are counted. Period at risk begins the later of January 1 or waiting list registration and ends on the earlier of December 31, date of death, or date of removal for other reasons. Please see Technical Notes for further details about death rate computation.

^{(*) =} For cohorts with fewer than 10 patients, rates are imprecise and are not reported.